

AD-A040 983

NAVAL RESEARCH LAB WASHINGTON D C
SPECTROSCOPIC OBSERVATION OF FAST IONS FROM LASER-PRODUCED PLAS--ETC(U)
MAY 77 E A MCLEAN, R DECOSTE, B H RIPIN
NRL-MR-3526

F/G 7/4

UNCLASSIFIED

NL

1 OF 1
AD
A040983



END

DATE
FILMED
7-77

ADA 040983

12
B.S.

NRL Memorandum Report 3526

Spectroscopic Observation of Fast Ions from Laser-Produced Plasmas


E. A. McLEAN, B. H. RIPIN, J. A. STAMPER, J. M. McMAHON and S. E. BODNER

*Laser Plasma Branch
Plasma Physics Division*

R. DECOSTE and H. R. GRIEM

*University of Maryland
College Park, Maryland*

May 1977



NAVAL RESEARCH LABORATORY
Washington, D.C.

AD No. _____
DDC FILE COPY

Approved for public release; distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)


REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 3526	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) SPECTROSCOPIC OBSERVATION OF FAST IONS FROM LASER-PRODUCED PLASMAS		5. TYPE OF REPORT & PERIOD COVERED Interim report on a continuing NRL problem.
7. AUTHOR(s) E.A. McLean, R. Decoste,* B.H. Ripin, J.A. Stamper, H.R. Griem,* J.M. McMahon and S.E. Bodner		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D.C. 20375		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS Energy Research and Development Administration Washington, D.C. 20545		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem H02-29A
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE May 1977
		13. NUMBER OF PAGES 10
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES *University of Maryland, College Park, Maryland. This work was performed at the Naval Research Laboratory under the auspices of the U.S. Energy Research and Development Administration.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Fast ions Nd:glass lasers Laser-plasma spectroscopy Ion energies Carbon ions		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Using a time-of-flight spectroscopic technique, measurements were made of the ion energy dis- tributions of very fast ions and thermal ions produced when a 7-15 J, 100 psec Nd:glass laser pulse (1.06 μm) strikes a $(\text{CH}_2)_n$ slab target. Ion energies greater than 0.5 MeV have been observed for the first time with this technique of measurement. A simultaneous comparison is made between the signal of an ion charge collector placed 30 cm from the target and the intensity of the C VI 3434 A ion line at 1 cm from the target. micrometer		


DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)





SPECTROSCOPIC OBSERVATION OF FAST IONS FROM LASER-PRODUCED PLASMAS

Exceedingly energetic ions ($v > 10^8$ cm sec $^{-1}$) are observed when high irradiance laser beams strike solid targets.¹ It has been established in these experiments that a large fraction of the absorbed laser energy is emitted in the form of kinetic energy of ions.^{2,3} The usual technique to measure the ion energy distribution is to use ion charge collectors, or an electrostatic or magnetic ion analyzer. In this paper a spectroscopic technique⁴⁻⁷ is described that allows measurements of the velocity distribution of the ions of a particular species of ionization. Further advantages of the technique are that simultaneous measurements can be made of both the high energy ions as well as the thermal ions and that measurements can be made close to the target (1-20 mm) and within the lens cone, without interfering with the incident laser beam. Typically, the spectral line intensities can be observed out to several centimeters from the target surface. Here we report the first observations of fast ions with energies beyond ~ 0.5 MeV using this optical technique. Previous studies⁴ using this technique, mostly at much lower incident laser power densities, did not show any ions with energies > 7 keV.

The excited ions are produced when a Nd:glass laser pulse (7-15 J, 100 psec) is focused with an f/1.9 aspheric lens onto a polyethylene (CH₂)_n slab target in an evacuated ($\sim 3 \times 10^{-4}$ Torr) chamber, as shown in Fig. 1. The half-energy focal spot diameter was 45 μ m, yielding an average focal irradiance of 10^{15} - 10^{16} W/cm². The plasma that is created is observed perpendicular to the laser axis by a 3/4-m monochromator and a fast photomultiplier (RCA-1P28). The

Note: Manuscript submitted May 13, 1977.

photomultiplier temporally resolves 5.5 \AA of the central intensity of a particular spectral line. The distance from the target to the point of observation and the time-of-flight of an ion bunch determines the mean velocity. Reflective optics were used to achieve an achromatic system which allows the optics to be aligned in the visible and then used down to about 2000 \AA . The position of observation could be varied by simply rotating the concave mirror with a spatial resolution of $\sim 0.1 \text{ mm}$. The photomultiplier was proved to be linear in the operating range, and the rise-time of the photomultiplier and recording electronics is better than 3 nsec . The time marker $t = 0$ was determined by observing, through the same optical system the initial rise of the $2\omega_0$ (5320 \AA) light generated at the target surface.

A time-integrated photographic spectrum ($2000 \text{ \AA} - 5000 \text{ \AA}$) showed many spectral lines of the carbon ions, C I - C VI. The continuum radiation was intense only within a few millimeters from the target surface and was below the noise level beyond 1 cm from the surface.

A typical oscillogram of the C VI 3434 \AA ($n = 6-7$) line is shown in Fig. 2. Here, the spectrograph was aligned to view the plasma 1 cm from the target. This trace shows low-intensity, but distinguishable, fast ion signals ($< 10 \text{ nsec}$) followed by the larger, more slowly rising peak due to the bulk plasma expansion (thermal ions). The first peak of the trace corresponds to an ion energy $\sim 560 \text{ keV}$, the second peak to an energy of $\sim 100 \text{ keV}$, and the last peak, or thermal peak, corresponds to an energy of $\sim 14 \text{ keV}$. The accuracy of the energy measurements of the fastest ions is about $\pm 50\%$ due to the comparable electronic and signal rise-times and the short time-of-flight. However, the accuracy of the thermal peak energy should be about $\pm 5\%$ since it occurs at a later time. The temporal accuracy of the fast ion measurement could be improved by making the measurement at 2 cm from the target surface, although the signal amplitude would decrease.

In Fig. 3 a comparison is made between the signal of an ion charge collector placed 30 cm from the target and 27° from the laser beam,

and the spectral intensity of the C VI 3434 \AA ion line at 1 cm from the target on the same shot. The target normal was rotated 10° with respect to the laser axis, and the laser beam was vertically polarized (s-polarization). The C VI curve has been normalized to the same velocity scale as the ion charge collector. Other than the initial spikes in the ion charge collector curve, the two curves are quite similar. For example, there are peaks on both curves corresponding to energies of $\sim 560 \text{ keV}$, $\sim 100 \text{ keV}$, and $\sim 15 \text{ keV}$. Other than minor differences in the energy at which the peaks occur, the largest difference between the two signals are in details in the structure of the ion charge collector signal. Other measurements using an electrostatic ion analyzer show that these narrow early peaks in the ion charge collector signal are mainly due to hydrogen ions.⁸ Also, we would not expect the amplitude of the two signals to track precisely since the secondary electron emission in the charge collector is dependent on the particle energy, and the line intensity is not simply related to the total number of ions. Although there are shot-to-shot variations in the fast ion traces, the general agreement between the ion charge collector and the C VI signals persisted for the shots taken under these conditions. Fast ion peaks were not seen in the oscillograms for carbon ions lower than C VI. This result is not fully understood at this time, but it may be that the fastest ions come from a very high temperature region where the ions are completely stripped and have just begun to pick up electrons on the way out. Recombination is evidently not fast enough to produce a significant population of lower charged states. Also, there is the possibility that we do not see fast ions in some of the lower stages of ionization because of differences in the radiative lifetimes of the transitions involved.⁹

In conclusion, this optical technique allows independent determinations of ion energies in laser-produced plasmas and unambiguously separates the various species of ionization. Some uncertainties encountered using the more common ion detectors are,

therefore, avoided. Measurements can be made less than 1 cm from the target surface and within the lens cone, where other ion detectors cannot easily be used. As has been pointed out by Boland, Irons, and McWhirter,⁴ it is possible using this technique and a separate measurement of the electron density and temperature to get absolute number densities of the various species of ions, and thus to determine the amount of energy in each type of ion and also the energy in the fast and thermal ions.

The authors are grateful for a useful discussion with Dr. R. W. P. McWhirter and for the technical assistance of E. C. Turbyfill, E. F. Vincent and T. H. DeRieux.

REFERENCES

1. A. W. Ehler, J. Appl. Phys. 46, 2464 (1975); I. Pelah, E. B. Goldman, and B. Yaakobi, Phys. Rev. Lett. 37, 829 (1976); V. C. Rupert and S. C. Gunn in Laser Program Annual Report - 1975, Lawrence Livermore Laboratory Report UCRL-50021-75, pp. 479-482; B. Luther-Davies and J. L. Hughes, Optics Comm. 18, 351 (1976); G. Charatis, J. Downward, R. Goforth, B. Guscott, T. Henderson, S. Hildum, R. Johnson, K. Moncur, T. Leonard, F. Mayer, S. Segall, L. Siebert, D. Solomon and C. Thomas, in Plasma Physics and Controlled Nuclear Fusion Research 1974, (IAEA, Vienna, 1975), Vol. II, p. 317.
2. Annual Progress Report on Laser Fusion Program, Institute of Laser Engineering, Osaka University, Suita, Osaka, Japan, Report ILE-APR-76 (Nov. 1976), p. 3-4 and 56-59.
3. J. Martineau, M. Rabeau, J. L. Bocher, J. P. Elie, and C. Patou, Optics Comm. 18, 347 (1976).
4. B. C. Boland, F. E. Irons, and R. W. P. McWhirter, J. Phys. B 1, 1180 (1968).
5. F. E. Irons, R. W. P. McWhirter and N. J. Peacock, J. Phys. B 5, 1975 (1972).
6. E. A. McLean, B. H. Ripin, J. A. Stamper, H. R. Griem, and S. E. Bodner, Bull. Am. Phys. Soc. 20, 1318 (1975); E. A. McLean, R. Decoste, B. H. Ripin, J. A. Stamper, H. R. Griem, and S. E. Bodner, Bull. Am. Phys. Soc. 21, 1084 (1976).
7. W. Seka, J. L. Schwob and C. Breton, J. Appl. Phys. 8, 3440 (1970).
8. R. Decoste and B. H. Ripin, submitted to Appl. Phys. Lett.
9. W. L. Wiese, M. W. Smith, and B. M. Glennon, Atomic Transition Probabilities, Vol. 1, (National Bureau of Standards, Wash., D.C., 1966).




Fig. 1 — Experimental arrangement




Fig. 2 — Oscillogram of the C VI 3434 Å ion line intensity observed 1 cm from the target surface. The time $t = 0$ is the time the laser pulse (7 J, 100 psec) initially strikes the target.




Fig. 3 — A comparison of the time trace of an ion charge collector and the intensity of the C VI 3434 Å ion line for the same laser shot. Both curves are normalized to the same energy and velocity scales. On the ion charge collector curve the spike adjacent to the left ordinate is due to x-ray flash and the next very sharp spike may be due to a spurious electromagnetic noise signal.